




November 22, 2011

**NEW DEGRADATION PRODUCT OF HBCD****Pentabromocyclododecene**

The widespread use of hexabromocyclododecane (HBCD) as an additive flame retardant for polystyrene foams and textiles has resulted in its detection in a variety of matrices. Unfortunately, the environmental impact of HBCD extends much further. HBCD has been shown to form decomposition products and metabolites through multiple transformation pathways such as debromination, dehydrobromination, and hydroxylation. As a result, the detection of HBCD related compounds, such as diastereomers of pentabromocyclododecene (PBCD), in environmental samples is on the rise. To date, PBCD has been detected in dust, eggs, fish, and sediments. Depending on the matrix and source, elevated levels of PBCD may be the result of bioaccumulation, due to its presence as a trace impurity in technical HBCD, or metabolism.

The main diastereomer formed from the decomposition of gamma HBCD is *rac*-(1,5*R*,6*S*,9*S*,10*R*)-pentabromocyclododecene. In order to aid researchers in the detection and identification of pentabromocyclododecene (PBCD) in environmental samples, Wellington now offers a reference standard of this single PBCD diastereomer.



Pentabromocyclododecene (PBCD)

Catalogue Number	Product (toluene)	Qty	Conc
PBCD	<i>rac</i> -(1,5 <i>R</i> ,6 <i>S</i> ,9 <i>S</i> ,10 <i>R</i> )-pentabromocyclododecene	1.2 ml	50 µg/ml

Please contact your local distributor or [info@well-labs.com](mailto:info@well-labs.com) for pricing and delivery.

Visit our website ([www.well-labs.com](http://www.well-labs.com)) for a complete listing of our new products.



ISO 9001

WELLINGTON LABORATORIES



Campro Scientific GmbH - Germany

P.O. Box 37 03 31 • 14133 Berlin • Phone +49.(0)30.629.01.89.0 • Fax +49.(0)30.629.01.89.89 • [info@campro.eu](mailto:info@campro.eu) • [www.campro.eu](http://www.campro.eu)